



HAL
open science

MORPHISMS FROM $S^2(\mathbb{C})$ TO $Gr(2, \mathbb{C}^4)$

Abdelghani El Mazouni, Fatima Laytimi, D.S. Nagaraj

► **To cite this version:**

Abdelghani El Mazouni, Fatima Laytimi, D.S. Nagaraj. MORPHISMS FROM $S^2(\mathbb{C})$ TO $Gr(2, \mathbb{C}^4)$. 2010. hal-00436261

HAL Id: hal-00436261

<https://univ-artois.hal.science/hal-00436261>

Preprint submitted on 5 Feb 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

MORPHISMS FROM $S^2(C)$ TO $Gr(2, \mathbb{C}^4)$.

A. EL MAZOUNI, F. LAYTIMI, AND D.S. NAGARAJ

ABSTRACT. In this note we study the natural vector bundle of rank 2 on $S^2(C)$, where $S^2(C)$ is the second symmetric power of non-singular irreducible non-degenerate space curve C .

Keywords: Vector bundles; Morphisms; Grassmannians; second symmetric power; curves.

1. INTRODUCTION

We denote by \mathbb{P}^3 the projective 3-space over the field \mathbb{C} of complex numbers and $Gr(2, \mathbb{C}^4)$ be Grassmannian of two-dimensional quotient spaces of \mathbb{C}^4 . Let $\mathcal{O}_{\mathbb{P}^3}(1)$ be the ample generator of the $\text{Pic}(\mathbb{P}^3)$. Let C be an irreducible non-singular curve in \mathbb{P}^3 . Assume C is non-degenerate, i.e., it is not contained in hyper plane. Let $L = \mathcal{O}_{\mathbb{P}^3}(1)|_C$. Let E_L denote the secant bundle of L , which is a rank 2 vector bundle on the second symmetric power $S^2(C)$ of the curve C . The bundle E_L is generated by 4 linearly independent sections. Hence there is a morphism $S^2(C)$ to $Gr(2, \mathbb{C}^4)$. In this article we study this morphism. Also, we study the properties of the restriction of E_L to some naturally imbedded curves in $S^2(C)$. Such a study was carried out by Schwarzenberger ([2]) when the curve C is a rational normal curve in \mathbb{P}^3 .

Acknowledgments: The third author would like to thank Université Lille and Université d'Artois, France.

2. SECANT BUNDLES

Let C be a smooth irreducible curve of genus g . By $S^2(C)$ we denote the second symmetric power of C , i.e., $S^2(C)$ is the quotient of $C \times C$ by the involution $(x, y) \mapsto (y, x)$, and $S^2(C)$ is smooth irreducible surface. A point of $S^2(C)$ can be thought of an effective divisor of degree 2 on C and $S^2(C)$ can be identified with the set of all effective divisor of degree two on C . Set

$$\Delta = \{(D, p) \in S^2(C) \times C \mid D = p + q\}.$$

1991 *Mathematics Subject Classification.* 14F17.

Then Δ is a divisor on $S^2(C) \times C$ which is known as the universal divisor of degree two and the first projection induces a surjective morphism

$$p_1 : \Delta \rightarrow S^2(C).$$

This is a two sheeted ramified covering. Let

$$p_2 : S^2(C) \times C \rightarrow C$$

be the second projection. For any line bundle M on C denote by E_M the locally free sheaf $(p_1)_*(p_2^*(M)|_\Delta)$ of rank two on $S^2(C)$. The exact sequence of $\mathcal{O}_{S^2(C) \times C}$ modules

$$(1) \quad 0 \rightarrow p_2^*(M)(-\Delta) \rightarrow p_2^*(M) \rightarrow p_2^*(M)|_\Delta \rightarrow 0$$

gives an exact sequence of $\mathcal{O}_{S^2(C)}$ modules

$$(2) \quad 0 \rightarrow p_{1*}(p_2^*(M)(-\Delta)) \rightarrow p_{1*}(p_2^*(M)) \rightarrow E_M.$$

By base change formula $p_{1*}(p_2^*(M)) = H^0(C, M) \otimes \mathcal{O}_{S^2(C)}$. If there is a subspace V of $H^0(C, M)$ such that the linear system given by V is very ample on C then we see that the induced map

$$V \otimes \mathcal{O}_{S^2(C)} \rightarrow E_M$$

of vector bundles is surjective. This defines a morphism

$$\phi : S^2(C) \rightarrow \text{Gr}(2, V).$$

If $\dim(V) = 2$ implies $\text{Gr}(2, V)$ is a point and if $\dim(V) = 3$ then $\text{Gr}(2, V) \simeq \mathbb{P}^2$ the morphism ϕ is finite and is a covering of degree $\binom{d}{2}$, where d is the degree of C in \mathbb{P}^2 . Hence the first interesting case is the case $\dim(V) = 4$, in that case as $\mathbb{P}(V) \simeq \mathbb{P}^3$. In the next section we treat this case.

Remark 2.1. *Let C be a non-singular irreducible curve of genus g and M be a line bundle of degree d . From the exact sequence (1) and Grothendick-Reimann-Rock one can compute the Chern classes of E_M :*

$$C_1(E_M) = (d - g - 1)x + \theta \quad \text{and} \quad C_2(E_M) = \binom{d - g}{2}x^2 + (d - g)x.\theta + \frac{\theta}{2}$$

where x is the cohomology class of image of $x \times C$ in $S^2(C)$ and θ is the cohomology class of the pull back of a theta divisor in $\text{Pic}^2(C)$ under the natural map of $S^2(C)$ into $\text{Pic}^2(C)$ [See, ACGH]. Note that the cohomology group $H^4(S^2(C), \mathbb{Z})$ is naturally isomorphic to \mathbb{Z} and

$$C_2(E_M) = \frac{(d - g)(d - g - 1)}{2} + (d - g)g + \frac{g(g - 1)}{2} = \frac{d^2 - d(2g + 1)}{2} + d.g.$$

3. SECANT BUNDLE ASSOCIATED TO A SPACE CURVE

Let C be a irreducible non-singular curve in \mathbb{P}^3 . Assume C is non-degenerate, i.e., it is not contained in hyper plane. Let $L = \mathcal{O}_{\mathbb{P}^3}(1)|_C$. Let E_L denote the secant bundle of L , which is a rank 2 vector bundle on the second symmetric power $S^2(C)$ of the curve C . Let V be the sub linear system of $H^0(C, L)$ given by the imbedding of C in \mathbb{P}^3 . As seen in the previous section the surjection of vector bundles

$$V \otimes \mathcal{O}_{S^2(C)} \rightarrow E_L$$

defines a morphism

$$\phi : S^2(C) \rightarrow \text{Gr}(2, V).$$

First we prove the following well known result which we include here due the lack of a good reference.

Theorem 3.1. *With the above notation the morphism ϕ is an imbedding then C is one of the following*

- 1) C is a rational normal curve of degree three
- 2) C is a smooth elliptic curve of degree four.

Proof: The curve C is non-degenerate implies the degree of the curve is greater than or equal to three. First we show that if the degree of C in \mathbb{P}^3 three or a smooth elliptic curve of degree four then it is an imbedding.

Case a): Assume degree of C is three. Then by projecting it from a point on it to a plane not containing it we see that C is isomorphic to \mathbb{P}^1 and the imbedding of C in \mathbb{P}^3 is given by the complete linear system $H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(3))$. In this case $S^2(C)$ is isomorphic to \mathbb{P}^2 and the line bundle $\mathcal{O}_{\mathbb{P}^2 \times \mathbb{P}^1}(\Delta)$ of the universal divisor Δ is isomorphic to $p_1^*(\mathcal{O}_{\mathbb{P}^2}(1)) \otimes p_2^*(\mathcal{O}_{\mathbb{P}^1}(2))$. Hence the exact sequence (2) becomes

$$(3) \quad 0 \rightarrow \mathcal{O}_{\mathbb{P}^2}(-1) \otimes H^0(\mathcal{O}_{\mathbb{P}^1}(1)) \rightarrow \mathcal{O}_{\mathbb{P}^2} \otimes H^0(\mathcal{O}_{\mathbb{P}^1}(3)) \rightarrow E_{\mathcal{O}_{\mathbb{P}^1}(3)} \rightarrow 0.$$

From the exact sequence (3) we see that the line bundle $\det(E_{\mathcal{O}_{\mathbb{P}^1}(3)})$ is isomorphic to $\mathcal{O}_{\mathbb{P}^2}(2)$ and the induced map

$$\overset{2}{\wedge} H^0(\mathcal{O}_{\mathbb{P}^1}(3)) \rightarrow H^0(\det(E_{\mathcal{O}_{\mathbb{P}^1}(3)}))$$

is an isomorphism. Since $\mathcal{O}_{\mathbb{P}^2}(2)$ gives an imbedding of \mathbb{P}^2 in $\mathbb{P}(H^0(\mathcal{O}_{\mathbb{P}^2}(2)))$ and $\text{Gr}(2, H^0(\mathcal{O}_{\mathbb{P}^1}(3)))$ is naturally imbedded in $\mathbb{P}(\overset{2}{\wedge} H^0(\mathcal{O}_{\mathbb{P}^1}(3)))$ we see that ϕ followed by the natural imbedding of $\text{Gr}(2, H^0(\mathcal{O}_{\mathbb{P}^1}(3)))$ in $\mathbb{P}(\overset{2}{\wedge} H^0(\mathcal{O}_{\mathbb{P}^1}(3)))$ is an imbedding. Hence ϕ is is an imbedding in this case.

case 2): Assume C in \mathbb{P}^3 is a nonsingular elliptic curve of degree four. Since degree of the line bundle L is four we see that by Riemann-Roch theorem $V = H^0(L)$ and $\dim(H^0(L^2))$ is eight. Hence there is a pencil of quadrics containing C . Since C is non-degenerate we see that all the quadrics in this pencil are irreducible and C is the scheme theoretic intersection of any two distinct members of this pencil. From this we can deduce that any line in \mathbb{P}^3 can meet C in at most two points. Also, if $Z \subset C$ is a closed sub-scheme supported on a finite set such that the vector space dimension $\ell(\mathcal{O}_Z)$ of \mathcal{O}_Z is three, then the natural map

$$H^0(L) \rightarrow \mathcal{O}_Z$$

is surjective. Hence we see that the morphism

$$\phi : S^2(C) \rightarrow \text{Gr}(2, V)$$

is an imbedding.

If C is a non degenerate non singular irreducible curve of degree four then C is either a rational curve or an elliptic curve. This can be seen by projecting C from a point on it to a plane not containing it. Moreover the curve C is a rational curve if and only if the projected curve is singular curve. Hence the curve C is rational if and only if there is line ℓ_0 in \mathbb{P}^3 such that the length of the scheme $C \cap \ell_0$ is greater or equal to three. Since the scheme $C \cap \ell_0$ determines a sub scheme of length greater or equal to two on $S^2(C)$ we see that

$$\phi : S^2(C) \rightarrow \text{Gr}(2, V)$$

is not an imbedding.

Let C be a irreducible non-singular curve in \mathbb{P}^3 of degree greater or equal to five. Then by using Castelnuov bound on the genus of a curve in terms of its degree ([4]) we see that there are lines in \mathbb{P}^3 which intersect C in a sub scheme of length greater or equal to three. Hence in this case

$$\phi : S^2(C) \rightarrow \text{Gr}(2, V)$$

is not an imbedding. This completes the proof of the theorem. \square

Remark 3.2. *Let C be a smooth curve in \mathbb{P}^3 of degree d greater than or equal to five. Then by Using Reimann-Roch theorem we see that projection h of C from a point p on C to a \mathbb{P}^2 not containing p is not an imbedding because*

$$h^*(\mathcal{O}_{\mathbb{P}^2}(1)) = \mathcal{O}_{\mathbb{P}^3}(1)|_C(-p).$$

From this we get through every point of $C \in \mathbb{P}^3$ there are lines which intersect C in more than two points (counting with multiplicity).

Remark 3.3. *On the Grassmaian $Gr(2, \mathbb{C}^4)$ one has universal exact sequence:*

$$(4) \quad 0 \rightarrow S \rightarrow \mathbb{C}^4 \otimes \mathcal{O}_{Gr(2, \mathbb{C}^4)} \rightarrow Q \rightarrow 0,$$

where S and Q are respectively the universal sub bundle and quotient bundle of rank two on $Gr(2, \mathbb{C}^4)$. The fiber of Q (resp. of S) at a point $p \in Gr(2, \mathbb{C}^4)$ is the two dimensional quotient space (resp. subspace, which is the kernal of this quotient map) of \mathbb{C}^4 corresponding to the point p . It is known (see, page 197 and 411 of [4]) that the cohomology group $H^4(Gr(2, \mathbb{C}^4), \mathbb{Z})$ is equal to

$$(5) \quad \mathbb{Z}[c_2(Q)] \oplus \mathbb{Z}[c_2(S)],$$

where $c_2(Q)$ (resp. $c_2(S)$) is the second Chern class of Q (resp. of S). The morphism $\phi : S^2(C) \rightarrow Gr(2, \mathbb{C}^4)$ determines a cohomology class in $H^4(Gr(2, \mathbb{C}^4), \mathbb{Z})$ namely $\phi_*([S^2(C)])$. By the identification ((5)) the cohomology class $\phi_*([S^2(C)])$ is represented by a pair of integers (a, b) . Using the geometry of $Gr(2, \mathbb{C}^4)$ we see that a is the number secants lines to C which passes through a generic point of \mathbb{P}^3 and b is the number of secant lines to C which are contained in generic plane of \mathbb{P}^3 (see [4]). From this description we see that if C is a rational normal curve of degree three than the cohomology class $\phi_*([S^2(C)])$ is $(1, 3)$ and if C is an elliptic curve of degree four than the cohomology class $\phi_*([S^2(C)])$ is $(2, 6)$.

4. PROPERTIES OF SECANT BUNDLE

Let C, L and V be as in previous section. When C is a rational normal curve in \mathbb{P}^3 from the results obtained by Schwarzenberger in ([2]) it follows that restriction of E_L is semi-stable for all lines in $S^2(C)$ which not tangent to the image of the diagonal of $C \times C$ and not semi-stable to those lines which are tangent to the image of the diagonal of $C \times C$. For a general curve C we have the following:

Theorem 4.1. *Let C be a non-singular curve of degree $d \geq 4$ in \mathbb{P}^3 and $L = \mathcal{O}_{\mathbb{P}^3}(1)|_C$. Then $E_L|_{x \times C}$ is not semi-stable for every $x \in C$.*

Proof: If $D = p + q$ be a point on the surface $S^2(C)$ then the fiber $E_L|_D$ of the vector bundle E_L at D can be identified with $L \otimes \mathcal{O}_D$, where \mathcal{O}_D is the structure sheaf sub-scheme of C of length two corresponding to D . Since every point of the curve $p \times C$ is of the form $p \times q$ for some $q \in C$ by looking at the linear system $V(-p) = \{s \in V | s(p) = 0\}$ we see that there is an injective homomorphism

$$0 \rightarrow L(-p) \rightarrow E_L|_{p \times C}.$$

But by (2.1) we see that $C_1(E_L|_{p \times C}) = d - 1$, where d is the degree of the line bundle L on C . Hence there exists a line bundle M of degree 0 on C such that $E_L|_{p \times C}$ fits in an exact sequence

$$0 \rightarrow L(-p) \rightarrow E_L|_{p \times C} \rightarrow M \rightarrow 0.$$

This proves that $E_L|_{p \times C}$ is not semi-stable as degree of M is less than half the degree of $E_L|_{p \times C}$.

REFERENCES

- [1] E. Arbarello, M. Cornalba, P. Griffiths and J. Harris: *Geometry of Algebraic curves I* Grundle. der Math. W. 267, Berlin-Hiedelberg-New York 1985.
- [2] R. L. E. Schwarzenberger: *Vector bundles on the projective plane* Proc. London Math. Soc. (3) 11 (1961) 623-40.
- [3] J. D’Almeida: *Une involution sur un espace de modules de fibrés instantons* Bull. Soc. Math. Fr. **128** (2000), 577–584.
- [4] P. Griffiths and J. Harris: *Principles of Algebraic Geometry*. Wiley Interscience Publication. 1978.
- [5] S. P. Inamdar and D. S. Nagaraj: *Cycle class map and restriction of subvarieties* J. Ramanujan Math. Soc. **17**, No.2(2002) 85–91.
- [6] F. Laytimi and D. S. Nagaraj: *Vector bundles generated by sections and morphism to Grassmannian* (preprint)
- [7] C. Okonek, M. Schneider and H. Spindler: *Vector bundles on complex projective spaces*. Progress in Mathematics, No. 3, Birkhäuser, Boston, Mass., 1980.
- [8] Hiroshi Tango: *On (n-1)-dimensional projective spaces contained in the Grassmann variety Gr(n, 1)*. J. Math. Kyoto Univ. 14-3 (1974) 415-460.
- [9] Sheng-Li Tan and Eckart Viehweg: *A note on Cayley-Bacharach property for vector bundles* Complex analysis and Algebraic geometry. A volume in memory of Michael Schneider. Editors: Thomes Peternell and Frank-Olaf Schreyer. Walter de Gruyter & Co., Berlin, (2000) 361-373.

LABORATOIRE DE MATHÉMATIQUES DE LENS EA 2462 FACULTÉ DES SCIENCES
JEAN PERRIN RUE JEAN SOUVRAZ, SP18 F-62307 LENS CEDEX FRANCE
E-mail address: mazouni@euler.univ-artois.fr

F. L.: MATHÉMATIQUES - BÂT. M2, UNIVERSITÉ LILLE 1, F-59655 VILLENEUVE D’ASCQ CEDEX, FRANCE
E-mail address: fatima.laytimi@math.univ-lille1.fr

INSTITUTE OF MATHEMATICAL SCIENCES C.I.T. CAMPUS, TARAMANI, CHENNAI 600113, INDIA
E-mail address: dsn@imsc.res.in