



HAL
open science

A new tractable subclass of the rectangle algebra

Philippe Balbiani, Jean-François Condotta, Luis Fariñas del Cerro

► **To cite this version:**

Philippe Balbiani, Jean-François Condotta, Luis Fariñas del Cerro. A new tractable subclass of the rectangle algebra. Sixteenth International Joint Conference on Artificial Intelligence (IJCAI 1999), The International Joint Conferences on Artificial Intelligence, Inc., 1999, Stockholm, Sweden. pp.442-447. hal-03300314

HAL Id: hal-03300314

<https://univ-artois.hal.science/hal-03300314>

Submitted on 2 Sep 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A new tractable subclass of the rectangle algebra

P. Balbiani

Laboratoire d'informatique de Paris-Nord
Avenue J.-B. Clement
F-93430 Villetaneuse, France
balbiani@lipn.univ-paris13.fr

J.-F. Condotta, L. Farinas del Cerro

Institut de recherche en informatique de Toulouse
1.18 route de Narbonne
F-31062 Toulouse Cedex 4, France
{condotta, farinas}@irit.fr

Abstract

This paper presents the 169 permitted relations between two rectangles whose sides are parallel to the axes of some orthogonal basis in a 2-dimensional Euclidean space. Elaborating rectangle algebra just like interval algebra, it defines the concept of convexity as well as the ones of weak preconvexity and strong preconvexity. It introduces afterwards the fundamental operations of intersection, composition and inversion and demonstrates that the concept of weak preconvexity is preserved by the operation of composition whereas the concept of strong preconvexity is preserved by the operation of intersection. Finally, fitting the propagation techniques conceived to solve interval networks, it shows that the polynomial path-consistency algorithm is a decision method for the problem of proving the consistency of strongly preconvex rectangle networks.

1 Introduction

Spatial representation and reasoning concern many areas of artificial intelligence: computer vision, geographic information, natural language understanding, computer-aided design, mental imagery, etc. These last ten years, numerous formalisms for reasoning about space were proposed [Egenhofer *et al*, 1991; Mukerjee *et al*, 1990; Freska, 1992; Randell *et al*, 1992]. We can mention as an example the well-known model of the regions proposed by Cohn, Cui and Randell [Randell *et al*, 1992] whose objects are the regions of a topological space and relations are eight topological relations, and for which Renz and Nebel [Renz *et al*, 1997] characterize a maximal tractable subclass of relations. Although this formalism is very attractive, it suffers from the impossibility of expressing orientation relations.

An example which enables this is the rectangle algebra (RA) [Giisgen, 1989; Mukerjee *et al*, 1990; Balbiani *et al*, 1998]. It is an extension for the space of the better known model for reasoning about time: the interval algebra (IA) proposed by Allen [Allen, 1983]. The basic objects of this spatial formalism are the rational rectangles

whose sides are parallel to the axes of some orthogonal basis in a 2-dimensional Euclidean space. Though restrictive, it is sufficient for applications in domains like architecture or geographic information. The relations between these objects are the 13 x 13 pairs of atomic relations which can hold between two rational intervals. These relations are very expressive, with them we can express both directional relations such as *left-of*, *right-of above*, etc., and topological relations such as *disjoint*, *overlap*, etc., between two rectangles.

In RA, spatial information are represented by spatial constraint networks which are special constraint satisfaction problems (CSPs). In these CSPs, each variable represents a rational rectangle and each constraint is represented by a relation of RA.

Given a spatial constraint network, the main problem is to know whether or not it is consistent, i.e, whether or not the spatial information represented by the network is coherent. Generally, this problem is NP-complete, but we can find subsets of the whole relations of RA for which this problem is polynomial, as in IA [Nebel *et al*, 1994; Beek, 1992]. Notably, in [Balbiani *et al*, 1998], Balbiani *et al*, presented a tractable set of relations called saturated-preconvex relations. In this paper, we present a new tractable set: the set of strongly-preconvex relations. This set contains the saturated-preconvex relations. We also prove that the well-known method, the path-consistency method, is complete for this set. To prove these results, we introduce a new method, called the weak path-consistency method, which is almost the path-consistency method.

The remainder of the paper is organised as follows. In Section 2, we make some recalls about RA, moreover we introduce the weakly-preconvex relations. Section 3 is concerned with some properties of fundamental operations: composition, intersection and inverse. In Section 4, we discuss spatial constraint network and we present the weak path-consistency method. Section 5 is concerned with the tractability results. In section 6, we define the strongly-preconvex relation and Section 7 concludes with suggestions for further extensions.

2 The rectangle algebra

The considered objects are the rectangles whose sides are parallel to the axes of some orthogonal basis in a 2-dimensional Euclidean space. The basic relations between these objects are defined from the basic relations of the interval algebra (denoted by \mathcal{B}_{int} and represented in the figure 2) in the following way :

$$\mathcal{B}_{rec} = \{(A, B) : A, B \in \mathcal{B}_{int}\}.$$

\mathcal{B}_{rec} constitutes the exhaustive list of the relations which can hold between two rational rectangles. For example, in figure 1, two rational rectangles satisfy the basic relation (m, p) of \mathcal{B}_{rec} . The set of the relations of the rect-




Figure 1: Two rational rectangles satisfying the basic relation (m, p) .

angle algebra is defined as the power set of \mathcal{B}_{rec} . Each relation of $2^{\mathcal{B}_{rec}}$ can be seen as the union of its basic relations. We obtain 2^{169} relations. Let R be a rectangle relation of $2^{\mathcal{B}_{rec}}$, we call the projections of R the two relations of $2^{\mathcal{B}_{int}}$, denoted by R_1 and R_2 and defined by:

$$R_1 = \{A : (A, B) \in R\} \text{ and } R_2 = \{B : (A, B) \in R\}.$$

Relation	Symbol	Reverse	Meaning	Dim
precedes	p	pi		2
meets	m	mi		1
overlaps	o	oi		2
starts	s	si		1
during	d	di		2
finishes	f	fi		1
equals	eq	eq		0

Figure 2: The set \mathcal{B}_{int} of the basic relations of 1A.

Ligozat [Ligozat, 1994] arranges the basic relations of the interval algebra in a partial order \leq which defines a




Figure 3: The interval lattice $(\mathcal{B}_{int}, \leq)$.

lattice called the interval lattice (see fig. 3). From this partial order we define the partial order \leq on \mathcal{B}_{rec} $(A, B) \leq (C, D)$ iff $A \leq C$ and $B \leq D$, with $(A, B), (C, D) \in \mathcal{B}_{rec}$.

$(\mathcal{B}_{rec}, \leq)$ defines a lattice that we call the rectangle lattice. The convex relations of IA and RA correspond to the intervals in the interval lattice and in the rectangle lattice respectively. We can prove that a convex relation R of RA is the cartesian product of its projections and that its projections are both convex relations of IA.

For each relation R of IA and RA, there exists a smallest, convex relation of 1A and RA respectively, which contains R . According to Ligozat's notation, we denote this relation by $I(R)$ (the convex closure of R). We can easily prove the following proposition:

Proposition 1 Let $R \in 2^{\mathcal{B}_{rec}}$ be, $I(R) = I(R_1) \times I(R_2)$.

In [Ligozat, 1994], a basic relation of 1A is represented by a region of the Euclidean plane and a relation of IA by the union of the regions representing its basic relations. With this representation two concepts are defined - the dimension and the topological closure of a relation of IA in the following way:

- the dimension of a basic relation A (see figure 2), denoted by $dim(A)$, corresponds to the dimension of its representation in the plane, i.e. 0, 1 or 2. The dimension of a relation R is the maximal dimension of its primitives.
- The topological closure of a basic relation A , denoted by $C(A)$ is the relation which corresponds to the topological closure of the region of A (see Table 1). For a relation R , $C(R)$ is the union of the topological closure of its basic relations.

These definitions use a geometrical representation of the basic relations, but we shall express them differently. In IA, each basic relation forces zero, one or two endpoint equalities. For example, between two intervals the basic relation eq imposes the first and the second endpoints to be equal whereas the basic relation p imposes no equality. We can use the following definition:

Definition 1 the dimension of a basic relation A of \mathcal{B}_{int} is the maximal number of equalities which can be imposed by a basic relation (i.e. 2) minus the number of equalities imposed by the basic relation A .

Relation	Topological closure	Relation	Topological closure
P	$\{p, m\}$	pi	$\{pi, mi\}$
m	$\{rn\}$	mi	$\{mi\}$
o	$\{rn, o, fi, s, eq\}$	oi	$\{mi, oi, f, si, eq\}$
s	$\{s, eq\}$	si	$\{si, eq\}$
d	$\{d, s, f, eq\}$	di	$\{rit, si, fi, eq\}$
f	$\{f, eq\}$	fi	$\{fi, eq\}$
eq	$\{eq\}$		

Table 1: The topological closure of the atomic relations of IA.

Then we can easily extend this definition to RA:

Definition 2 *the dimension of a basic relation of \mathcal{B}_{rec} is the maximal number of endpoint equalities that a basic relation can impose between the orthogonal axe projections of two rectangles (i.e. 4) minus the member of endpoint equalities imposed by the basic relation.*

For example, the dimension of the basic relation (m, p) (see figure 1) is 3. The dimension of a relation R of $2^{\mathcal{B}_{rec}}$ is the maximal dimension of its basic relations, and we suppose that the dimension of the empty relation is -1 . It is easy to prove the following facts:

Proposition 2

- let (A, B) be a basic relation of \mathcal{B}_{rec} , $dim((A, B)) = dim(A) + dim(B)$;
- let $R \in 2^{\mathcal{B}_{rec}}$ be a convex relation, $dim(R) = dim(R_1) + dim(R_2)$.

To extend the definition of the topological closure to RA we use directly the topological closure of the atomic relations of IA (see Table 1) by the following ways:

Definition 3

- let $(A, B) \in \mathcal{B}_{rec}$ be, $C((A, B)) = C(A) \times C(B)$;
- let $R \in 2^{\mathcal{B}_{rec}}$ be, $C(R) = \bigcup \{C((A, B)) : (A, B) \in R\}$.

Although speaking of topological closure doesn't, make any sense because we don't use regions to represent the relations of RA, we shall continue to use the term as in IA. We can prove the following proposition :

Proposition 3

- let $R, S \in 2^{\mathcal{B}_{rec}}$ be, $C(R) \circ C(S) \subseteq C(R \circ S)$;
- let R be a convex relation of $2^{\mathcal{B}_{rec}}$, $C(R) = C(R_1) \times C(R_2)$;
- let R be a convex relation of $2^{\mathcal{B}_{rec}}$, for all $(A, B) \in R$ there exists $(C, D) \in R$ such that $dim((C, D)) = dim(R)$ and $(A, B) \in C((C, D))$.

Proof.(Sketch) Ligozat proves the first part of this proposition for two relations of IA. Using this result we can also prove this for RA. The proof of the second part is obvious. To prove the third one, first we prove it for IA by examining the exhaustive list of the convex relations of IA. And from this result and the first two parts we prove the result for RA. \square

In [Ligozat, 1994], the preconvex relations of IA are defined in the following way: a relation R of IA is a preconvex relation iff $dim(I(R) \setminus R) < dim(R)$ – an equivalent definition is: R is a preconvex relation iff $I(R) \subseteq C(R)$. The set of preconvex relations coincides with the set of the well known ORD-Horn relations, which is the maximal tractable set of IA which contains all basic relations [Nebel et al., 1994]. Using the convex closure and the dimension of a relation of RA, now, we are ready to define the preconvex relations in RA. We shall call these relations the weakly-preconvex relations.

Definition 4 *Let $R \in 2^{\mathcal{B}_{rec}}$ be, R is a weakly-preconvex relation iff $dim(I(R) \setminus R) < dim(R)$.*

Using proposition 3, we can prove:

Proposition 4 *A relation R of RA is a weakly-preconvex relation iff $I(R) \subseteq C(R)$.*

From this proposition it follows that:

Proposition 5 *A relation R of RA is a weakly-preconvex relation iff $C(R)$ is a convex relation.*

Proof. Let R be a weakly-preconvex relation of RA. From proposition 4, $R \subseteq I(R) \subseteq C(R)$. So, $C(R) \subseteq C(I(R)) \subseteq C(C(R))$. Since $C(C(R)) = C(R)$, we deduce that $C(R) = C(I(R))$. Given that every convex relation of IA has a convex topological closure [Ligozat, 1996], from propositions 1 and 3 we deduce that $C(I(R))$ is convex. Let, $R \in 2^{\mathcal{B}_{rec}}$ be such that $C(R)$ is convex. Since $R \subseteq C(R)$, by definition of I we deduce that $I(R) \subseteq C(R)$. \square

3 Fundamental operations

The set $2^{\mathcal{B}_{rec}}$ is enriched with the fundamental relational operations, the binary operations, intersect (\cap), composition (\circ) and the unary operation inverse ($^{-1}$). The composition of two basic relations and the inverse of a basic relation in RA can be computed from the same operations in IA in the following way : $(A, B) \circ (C, D) = (A \circ C) \times (B \circ D)$ and $(A, B)^{-1} = (A^{-1}, B^{-1})$. Composition between two relations in $2^{\mathcal{B}_{rec}}$ is defined by : $R \circ S = \bigcup \{(A, B) \circ (C, D) : (A, B) \in R, (C, D) \in S\}$. The inverse of a relation in $2^{\mathcal{B}_{rec}}$ is the union of the inverse of its basic relations. $2^{\mathcal{B}_{rec}}$ is stable for these three operations. Considering a subset of $2^{\mathcal{B}_{rec}}$ firstly we must look whether this subset is stable for these operations. We can easily prove the following proposition:

Proposition 6 *for every relation $R, S, T, U \in 2^{\mathcal{B}_{rec}}$,*

- $(R \times S) \cap (T \times U) = (R \cap T) \times (S \cap U)$;
- $(R \times S) \circ (T \times U) = (R \circ T) \times (S \circ U)$;
- $(R \times S)^{-1} = R^{-1} \times S^{-1}$.

Consequently, the set of the convex relations of $2^{\mathcal{B}_{rec}}$ is stable for the operations: intersection, composition and inverse. Hence the set of the convex relations of $2^{\mathcal{B}_{rec}}$ is a subclass of RA. From all this it follows that:

Proposition 7 *For all relations R and $S \in 2^{\mathcal{B}_{rec}}$, $I(R \circ S) \subseteq I(R) \circ I(S)$.*

Indeed, $I(R) \circ I(S)$ is a convex relation and contains the relation $R \circ S$.

Now, let us consider the subset of the weakly-preconvex relations, we denote this set by \mathcal{W} . We have the following result:

Proposition 8 \mathcal{W} is stable for the operations \circ , $^{-1}$ but it is not stable for the operation \cap .

Proof. The proof of the stability of the composition is the same as the one given in [Ligozat, 1994] for the stability of the preconvex relations of IA: for all relations $R, S \in \mathcal{W}$, $I(R \circ S) \subseteq I(R) \circ I(S) \subseteq C(R) \circ C(S) \subseteq C(R \circ S)$. Hence, $R \circ S$ is a weakly-preconvex relation. We notice that for all basic relation $(A, B) \in \mathcal{B}_{rec}$, we have $C((A, B)^{-1}) = C((A, B))^{-1}$, so for all relation $R \in 2^{\mathcal{B}_{rec}}$, $C(R^{-1}) = C(R)^{-1}$. Moreover, owing to the symmetry of the rectangle lattice, we have $I(R^{-1}) = I(R)^{-1}$. From these results, the stability of \mathcal{W} for the operation inverse is obvious. \square

Unlike the case of IA, the weakly-preconvex relations in HA are not stable for the intersection. We can see it with the following counter-example. Let consider the two weakly-preconvex relations R, S of $2^{\mathcal{B}_{rec}}$ defined by $R = \{(o, o), (eq, s), (s, eq)\}$ and $S = \{(d, d), (eq, s), (s, eq)\}$. $R \cap S$ is the relation $\{(eq, s), (s, eq)\}$, this relation is not weakly-preconvex. This lack of stability - as we may see in the sequel - will raise some problem.

4 Constraints networks

We start, this section with some reminders about constraints networks of RA [Balbiani et al., 1998]. A rectangle network \mathcal{N} is a structure (V, C) , where $V = \{V_1, \dots, V_n\}$ is a set of variables which represent, rational rectangles, and C is a mapping from $V \times V$ to the set $2^{\mathcal{B}_{rec}}$ which represent the binary constraints between the rational rectangles. C is such that :

- for every $i \in \{1, \dots, |V|\}$, $C_{ii} = \{(eq, eq)\}$;
- for every $i, j \in \{1, \dots, |V|\}$, $C_{ij} = C_{ji}^{-1}$.

An interval network is defined in the same way except that BV represents a set of rational intervals and C is to $2^{\mathcal{B}_{int}}$.

Let $\mathcal{N} = (V, C)$ be a rectangle network, the consistency' and the path-consistency of \mathcal{N} is defined in the following way:

- \mathcal{N} is consistent iff there exists a mapping m from V to the set of rational rectangles such that, for every $i, j \in \{1, \dots, |V|\}$, there exists a basic relation P such that $P \in C_{ij}$ and m_i and m_j satisfy P . We call such a mapping m , a satisfying instantiation of the network \mathcal{N} .
- ♦ \mathcal{N} is path-consistent when, for every $i, j \in \{1, \dots, |V|\}$, $C_{ij} \neq \{\}$ and for every $i, j, k \in \{1, \dots, |V|\}$, $C_{ij} \subseteq C_{ik} \circ C_{kj}$.

We have the same definitions for the interval networks. A convex (respectively weakly-preconvex) network is a

network which contains only convex relations (respectively weakly-preconvex relations).

We define another property, the weak path-consistency. A weakly path-consistent network is an almost path-consistent network:

Definition 5 A rectangle: (or interval) network is weakly path-consistent when, for every $i, j \in \{1, \dots, |V|\}$, $C_{ij} \neq \{\}$ and for every $i, j, k \in \{1, \dots, |V|\}$, $C_{ij} \subseteq I(C_{ik} \circ C_{kj})$.

A path-consistent, network is obviously weakly path-consistent, the contrary is not true.

A well-known polynomial method to see whether a network $\mathcal{N} = (V, C)$ is consistent, is the path-consistency method [Allen, 1983]. It consists to iterate on the network the triangulation operation: $C_{ij} := C_{ij} \cap (C_{ik} \circ C_{kj})$ for all $i, j, k \in \{1, \dots, |V|\}$, until we obtain a stable network. It is sound but not complete as a decision procedure for the issue of the consistency of a network. Indeed, if after applying the method we obtain the empty relation then the network is not consistent, in the contrary case we do not know whether the network is consistent.

Similarly, we define a weaker method that we call the weak path-consistency method. It consists to iterate the weak triangulation operation $C_{ij} := C_{ij} \cap I(C_{ik} \circ C_{kj})$. The time complexity is the same as the one of the path-consistency method. Moreover, since $(C_{ik} \circ C_{kj}) \subseteq I(C_{ik} \circ C_{kj})$, this method is sound but not complete like the path-consistency method.

5 Results of tractability

Let $\mathcal{N} = (V, C)$ be a network and m be a satisfying instantiation of \mathcal{N} , we shall say that m is maximal iff for all $i, j \in \{1, \dots, |V|\}$, $dim(P) = dim(C_{ij})$, with P the basic relation satisfied between m_i and m_j .

Ligozat [Ligozat, 1994] proves this important theorem:

Theorem 1 Let \mathcal{N} be an interval network which contains only preconvex relations, if \mathcal{N} is path-consistent then there exists a maximal instantiation of \mathcal{N} .

From this theorem we can prove the following lemma :

Lemma 1 Let $\mathcal{N} = (V, C)$ be a convex rectangle network, if \mathcal{N} is path-consistent then there exists a maximal instantiation of \mathcal{N} .

Proof. Let $\mathcal{N}' = (V', C')$ and $\mathcal{N}'' = (V'', C'')$ be two interval networks defined by $V = V' = V''$ and for all $i, j \in \{1, \dots, |V|\}$, $C'_{ij} = (C_{ij})_1$ and $C''_{ij} = (C_{ij})_2$. We can prove that \mathcal{N}' and \mathcal{N}'' are path-consistent, see [Balbiani et al., 1998]. Moreover they are convex (a fortiori preconvex¹). So there exists a maximal instantiation of \mathcal{N}' and \mathcal{N}'' , let us call them m' and m'' respectively. Let us consider the instantiation m of \mathcal{N} such that for every $i, j \in \{1, \dots, |V|\}$, m'_{ij} is the projection of the rational rectangle m_{ij} onto the horizontal axis and m''_{ij} is the projection of m_{ij} onto the vertical axis.

¹ We remind that the set of the preconvex relations contains the set of the convex relations in IA.

It is easy to see that m is a satisfying instantiation of \mathcal{N} . Moreover, m'_i (respectively m''_i) and m'_j (respectively m''_j) satisfy a. basic relation A (respectively B) such that $\dim(A) = \dim(C'_{ij})$ (respectively $\dim(B) = \dim(C''_{ij})$). So m_i and m_j are such that $\dim((A, B)) = \dim(C_{ij})$ because C_{ij} is a convex relation. So, m is a maximal instantiation of \mathcal{N} . \square

Consequently, for the weakly-preconvex relations of $2^{\mathcal{B}^{rec}}$, we have the following result:

Lemma 2 *Let $xY = (V', C)$ be a weakly-preconvex rectangle network, if N is weakly path-consistent then N is consistent, moreover there exists a maximal instantiation of N .*

Proof. Let $\mathcal{N}' = (V', C')$ be the rectangle network such that $V' = V$ and for all $i, j \in \{1, \dots, |V|\}$, $C'_{ij} = I(C_{ij})$. First, let us prove that \mathcal{N}' is a path-consistent network: for all $i, j, k \in \{1, \dots, |V|\}$ we have: $C_{ij} \subseteq I(C_{ik} \circ C_{kj})$, then $I(C_{ij}) \subseteq I(I(C_{ik} \circ C_{kj}))$, consequently $I(C_{ij}) \subseteq I(C_{ik} \circ C_{kj})$ and because $I(R \circ S) \subseteq I(R) \circ I(S)$ for all relations $R, S \in 2^{\mathcal{B}^{rec}}$, we obtain the result $I(C_{ij}) \subseteq I(C_{ik}) \circ I(C_{kj})$. So, $C'_{ij} \subseteq C'_{ik} \circ C'_{kj}$, hence \mathcal{N}' is path-consistent.

\mathcal{N}' is convex and path-consistent then there exists a maximal instantiation m of \mathcal{N}' . For each $i, j \in \{1, \dots, |V|\}$, m_i and m_j satisfy a basic relation P of C'_{ij} such that $\dim(P) = C'_{ij}$. C_{ij} is weakly-preconvex then $\dim(I(C_{ij}) \setminus C_{ij}) < C_{ij}$. Moreover, remind that $C'_{ij} = I(C_{ij})$, consequently $P \in C_{ij}$ and $\dim(P) = \dim(C_{ij})$. Hence, m is also a maximal instantiation of the rectangle network \mathcal{N} . \square

Because a path-consistent network is also weakly path-consistent too, this lemma is true as well for a weakly-preconvex network \mathcal{N} which is path-consistent instead of weakly path-consistent.

When we apply the path-consistency method from a weakly-preconvex network, we are not sure to obtain a weakly-preconvex network, because the set of the weakly-preconvex relations is not stable for the operation intersection, likewise with weak path-consistency method because the intersection between a weakly-preconvex relation and a convex relation is not always a weakly-preconvex relation. For example, the intersection between the weakly-preconvex relation $\{(d, d), (eq, s), (s, eq)\}$ and the convex relation $\{(s, s), (eq, s), (s, eq), (eq, eq)\}$ is $\{(s, eq), (eq, s)\}$ which is not weakly-preconvex. Hence, despite the previous lemma, we cannot, assert that the path-consistency method or the weak path-consistency method are decision procedures for the issue of the consistency of a rectangle network which contains only weakly-preconvex relations. But we can characterise some subsets of $2^{\mathcal{B}^{rec}}$ for which it works:

Theorem 2 (Main result) *Let $E \subset \mathcal{W}$, stable for the intersection with the convex relations.*

The weak path-consistency method is a decision procedure for the issue of the consistency of a rectangle network which contains just constraints in E .

Proof. From the previous lemma the proof is direct. Let \mathcal{N} be a rectangle network whose constraints are in E . After applying the weak path-consistency method from \mathcal{N} , we obtain a network \mathcal{N}' whose constraints are always in E and consequently which is weakly-preconvex. Moreover, \mathcal{N}' is equivalent to \mathcal{N} , so if \mathcal{N}' contains the empty relation then \mathcal{N}' and \mathcal{N} are not consistent. Else, from the previous lemma we deduce that there exists a maximal instantiation of \mathcal{N}' and consequently \mathcal{N}' and \mathcal{N} are consistent. \square

Moreover, we can assert that path-consistency method is also complete for rectangle networks whose relations are in such a set E .

Let, us denote by \overline{F} the closure of a set F of relations by the operations intersection, composition and inverse. Nebel and Burckert [Nebel *et al.*, 1994] show that from an interval network whose constraints are in \overline{F} we can construct, in polynomial time, another interval network whose constraints are in F and such that the former network is consistent, if and only if the latter is consistent. We can prove the same thing in RA. Hence, if E is a set of rectangle relations having the properties of the previous theorem, then the problem of the consistency of the networks whose relations belong to \overline{E} is a polynomial problem.

6 The strongly-preconvex relations

In this section, first, we are going to define a new set with the properties of theorem 2: the set of the strongly-preconvex relations of HA. Then, we shall prove that this set is the maximal set for these properties.

The strongly-preconvex relations of RA are defined by:

Definition 6 *Let R be a relation of $2^{\mathcal{B}^{rec}}$, R is a strongly-preconvex relation if, and only if, for all convex relations S , $R \cap S$ is a weakly-preconvex relation.*

The universal relation of $2^{\mathcal{B}^{rec}}$ is a convex relation, from this we deduce that a strongly-preconvex relation is a weakly-preconvex relation. We denote the set of the strongly-preconvex relations by \mathcal{S} . The convex relations are stable for the intersection and are weakly-preconvex relations. So, they belong to \mathcal{S} . What is more:

Proposition 9 *If $R, S \in \mathcal{S}$ then $R \cap S \in \mathcal{W}$.*

Proof. First, we are going to prove that $I(C(R \cap S)) = C(R \cap S)$. Let us denote $I(C(R \cap S))$ by T .

We have $R \cap S \subseteq T$, so $R \cap S \subseteq T \cap R$ and $R \cap S \subseteq T \cap S$. Consequently $T \subseteq I(C(T \cap R))$ and $T \subseteq I(C(T \cap S))$. Because R and S are strongly-preconvex relations and T is a convex relation, $T \cap R$ and $T \cap S$ are two weakly-preconvex relations. Consequently $C(T \cap R)$ and $C(T \cap S)$ are two convex relations hence, $T \subseteq C(T \cap R)$ and $T \subseteq C(T \cap S)$. When we compute the topological closure of a relation, we add only basic relations of lower dimension. So, $\dim(T) = \dim(C(T \cap R))$ and $\dim(T) = \dim(C(T \cap S))$. Let P be a basic relation such $P \in T$ and $\dim(P) = \dim(T)$. From the previous results we deduce that $P \in C(T \cap R)$, $P \in C(T \cap S)$, and $\dim(P) = \dim(C(T \cap R)) = \dim(C(T \cap S))$. So, as $T \cap R$ and

$T \cap S$ are weakly-preconvex relations, $P \in T \cap R$ and $P \in T \cap S$, hence $P \in C(R \cap S)$. Let Q be any basic relation belonging to T from proposition 3, we can deduce too that $Q \in C(R \cap S)$. Hence $T \subseteq C(R \cap S)$.

$C(R \cap S) \subseteq I(C(R \cap S)) \subseteq C(R \cap S)$, so $R \cap S$ is a weakly-preconvex relation of $2^{\mathcal{B}_{rec}}$. \square

Moreover, we can prove the following result:

Proposition 10 S is stable for the intersection with the convex relations of $2^{\mathcal{B}_{rec}}$.

Proof. Let R and S be two relations of $2^{\mathcal{B}_{rec}}$ such that R is a strongly-preconvex relation and S a convex relation. Let us prove that $R \cap S$ is a strongly-preconvex relation. Let T be any convex relation, we have: $(R \cap S) \cap T = R \cap (S \cap T)$. $S \cap T$ is a convex relation too. R is a strongly-preconvex relation, we can deduce that $R \cap (S \cap T)$ is a weakly-preconvex relation, hence $R \cap S$ is a strongly-preconvex relation. \square

Consequently, we can apply theorem 2 on the set S . From the last two propositions we can easily prove that \mathcal{S} is stable for the intersection. S is stable for the inverse operation too. But we have not yet succeeded in proving the composition stability, so perhaps $\overline{\mathcal{S}} = S$. Moreover, let E be a set with the properties of theorem 2, we can easily prove that $E \subseteq \mathcal{S}$. From all this it follows:

Theorem 3 S is tractable and \mathcal{S} is the maximal set included in \mathcal{W} and stable for the intersection with the convex relations.

Hence, \mathcal{S} is the maximal set with the properties of theorem 2.

A saturated-preconvex relation of $2^{\mathcal{B}_{rec}}$ corresponds to the cartesian product of two preconvex relations of $2^{\mathcal{B}_{int}}$ [Balbiani et al., 1998]. By using this and the fact that the set of the saturated-preconvex relations is stable for the intersection [Balbiani et al. 1998], from proposition 3 we can prove that the set of the saturated-preconvex relations is a subset, of \mathcal{S} . Moreover, it is a proper subset of \mathcal{S} . For example, let us consider the relation $\{(o, o), (s, eq), (eq, s), (s, s)\}$ which is strongly-preconvex but not saturated-preconvex. So, now \mathcal{S} is the largest known set to be a tractable set which contains the basic relations. This result provides also another proof of the tractability of saturated-preconvex relations.

7 Conclusion

The subclass generated by the set of the strongly-preconvex relations is now the biggest known tractable set of RA which contains the 169 atomic relations. An open question is: is this subclass a maximal tractable subclass which contains the atomic relations?

Another future development, is to extend RA to a greater dimension than two. For dimension n ($n \geq 3$), the considered objects will be the blocks whose sides are parallel to the axes of some orthogonal basis in a n -dimensional Euclidean space. The atomic relations between these objects are the 13^n relations obtained by the cartesian product, of atomic relations of IA. A first attempt shows that, the previous tractability results of RA can be easily extended to this structure.

References

- [Allen, 1983] J. Allen. *Maintaining knowledge about temporal intervals*. Communications of the ACM, Volume 26, 832-843, 1983.
- [Balbiani et al., 1998] P. Balbiani, J-F. Condotta and L. Farinas del Cerro. *A model for reasoning about bidimensional temporal relations*. A.G. Colin, L. Schubert, and S.C. Shapiro (editors), KR/98. Proc of the 6th Int. Conf., 124-130, Morgan Kaufmann, 1998.
- [Egenhofer et al., 1991] M. Egenhofer and R. Franzosa. *Point-set topological spatial relations*. International Journal of Geographical Information Svstems 5(2), 161-174, 1991.
- [Freska, 1992] C. Preska. *Using orientation information for qualitative spatial reasoning*. A U. Frank, I. Campari, and U. Formentini (editors), Theories and Methods of Spatio Temporal Reasoning in Geographic Space, Proc. of the Int. Conf. CIS - From Space to Territory, 162-178, Springer Verlag, 1992.
- [Giisgen, 1989] H. Giisgen. *Spatial reasoning based on Aliens temporal logic*. Report 1CSI TR89-049, International Computer Science Institute, 1989.
- [Ligozat, 1994] G. Ligozat. *Tractable relations in temporal reasoning: pre-convex relations*. ECAI-94. Workshop on Spatial and Temporal Reasoning, 99-108, 1994.
- [Ligozat, 1996] G. Ligozat. *A new proof of tractability for ORD-Horn relations*. AAI-96. Proc. of the 13th Nat. Conf. on Artificial Intelligence, Volume one, 395-401, AAI Press and MIT Press, 1996.
- [Nebel et al., 1994] B. Nebel, IP J. Biirckert. *Reasoning about temporal relations: a maximal tractable subclass of Allen's interval algebra*. AAI 94. Proc. of the 12th Nat. Conf. on Artificial Intelligence. Volume one, 356-361, AAI Press and MIT Press, 1994.
- [Mukerjee et al., 1990] A. Mukerjee, G. Joe. *Qualitative Model For Space*. AAI-90. Proc. of the 8th Nat. Conf. on Artificial Intelligence, Volume two, 721-727, AAI Press and MIT Press, 1990.
- [Randell et al., 1992] 1). Randell, Z. Cui, T. Cohn. *An interval logic for space based on "connection"*. B. Neumann (editor), ECAI-92. Proc. of the 10th European Conf. on Artificial Intelligence, 394-398, Wiley, 1992.
- [Renz et al., 1997] J. Renz, B. Nebel. *On the complexity of qualitative spatial reasoning: a maximal tractable fragment of the region connection calculus*. M. Polack (editor), IJCAI-97. Proc. of the 15th Int. Joint Conf. on Artificial Intelligence, Volume one, 522-527, Morgan Kaufmann, 1997.
- [Beek, 1992] P. van Beek. *Reasoning about qualitative temporal information*. Artificial Intelligence, 58(1-3), 297-321, 1992.